







Research Article

Pre-Hispanic anthropogenic wetlands in the upper Ica drainage, south-central Andes: dating and context

Kevin Lane^{1,*} , David Beresford-Jones² , Luis Coll¹, Erik Marsh³ , Rob Scaife⁴, Catriel Greco⁵ , Oliver Huaman Oros⁶, Alexander Herrera⁷, Jennifer Grant⁸ & Charles French^{9,*}

¹ Consejo Nacional de Investigación de Ciencia y Tecnología (CONICET), Instituto de las Culturas, Universidad de Buenos Aires, Argentina

² Heinz Heinen Centre for Advanced Study, University of Bonn, Germany

³ CONICET, Instituto Interdisciplinario de Ciencias Básicas, Universidad Nacional de Cuyo, Mendoza, Argentina

⁴ Geography and Environmental Science, University of Southampton, UK

⁵ CONICET, Facultad de Ciencias Físico Matemáticas y Naturales, Universidad Nacional de San Luis, Argentina

⁶ Proyecto Qhapaq Ñan, Sede Nacional, Ministerio de Cultura de Perú, Lima, Peru

⁷ Universidad de los Andes, Departamento de Historia del Arte, Bogotá, Colombia

⁸ CONICET, Instituto Nacional de Antropología y Pensamiento Latinoamericano, Buenos Aires, Argentina

⁹ Department of Archaeology, University of Cambridge, UK

* Authors for correspondences ✉ kevin.lane@cantab.net & caif2@cam.ac.uk



Humans have engineered their environments throughout the Holocene, especially in the construction of hydraulic infrastructure. In many regions, however, this infrastructure is difficult to date, including the vestiges of water-management systems in the Andean highlands. Focusing on silt reservoirs in the upper Ica drainage, Peru, the authors use cores and radiocarbon dates to demonstrate the pre-Hispanic construction of walls to enhance and expand wetlands for camelid pasture. Interventions dated to the Inca period (AD 1400–1532) indicate an intensification of investment in hydraulic infrastructure to expand production capacity in support of the state. The results are discussed in the context of the hydraulic strategies of other states and empires.

Keywords: Peru, Ica drainage, Inca Empire, anthropogenic wetland, hydrological technology

Introduction

Humans have been altering the physical environment for over 12 000 years and the Andes are no exception (Denevan 1992; Ellis *et al.* 2021). In the seemingly bleak, *suní-puna*

Received: 6 July 2021; Revised: 27 October 2021; Accepted: 10 December 2021

© The Author(s), 2022. Published by Cambridge University Press on behalf of Antiquity Publications Ltd. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

alpine-tundra of the high Andes, pre-Hispanic populations used and developed existing natural moorlands and wetlands—the latter known as *bofedales* in the Central Andes or *vegas* in the Southern Andes (Lane 2006). Here, we examine how these pre-Hispanic human populations created or augmented bofedales. This was usually achieved through the construction of canals or walled silt-traps in order to improve water and soil retention, thus providing more ample and better-quality pasturage. Dating such structures, especially silt-traps, however, is fraught with difficulty, “given [the] disturbance by erosion and deposition, by the process of agriculture, and by frequent rebuilding” (Denevan 2001: 172) that characterises their construction and use.

The results from an ongoing research project in the upper Ica drainage in southern Peru have revealed a series of walled silt traps or check dam-type structures, known as silt reservoirs. These were built to withhold silt and water, leading to peat and wetland growth at the summit of the Quebrada Marccaranca, where it opens into a natural basin. At some point in the past, the construction of these silt reservoirs led to the anthropogenic alteration of this natural basin, increasing the size of the wetlands. A series of three borehole samples and six radiocarbon dates from the basin’s peat levels reveal that this anthropogenic alteration probably occurred during the Late Horizon or Inca period (AD 1400–1532), before the area’s abandonment by the end of the eighteenth century.

In this article, we describe the use of silt reservoirs in the Andes, focusing particularly on the highest reaches of the upper Ica drainage, before charting the probable history of construction and use of these structures in the Quebrada Marccaranca. Finally, we consider the reasons for investment in these hydrological structures during the Inca period, analysing potential issues of imperial management of resources, land and surplus.

Anthropogenic wetlands in the Andes

Bofedales are poorly drained, high-altitude peatlands replete with cushion-like plants, such as *Distichia muscooides*, which, in turn, permit the growth of high-quality, softer grasses, including *Calamagrostis corymbosa* and *Calamagrostis rigescens* (Squeo *et al.* 2006; Maldonado Fonkén 2014). These wetlands and their rich plant communities are found almost exclusively within the *puna* ecozone—an alpine-tundra grassland located above 4000m asl in the Central Andes. Bofedales provide an excellent herding habitat that is highly coveted by modern-day pastoralist communities (Browman 1990); this was likely the case in the past, as well.

The *puna* is considered a critical resource for Andean highland communities, constituting 30 per cent of the Central Andean landscape; within this, bofedales constitute 2.5 per cent of the *puna* (Ravines 1978; Kuznar 1995: 36; Young *et al.* 1997). Even though bofedales provide optimum pasture for grazing, their use for herding is seasonal and generally restricted to the dry months just before the rains—usually between June and September. At other times of the year, the natural *sunipuna* pasture is normally sufficient for the needs of domesticated animals.

Given that the importance of bofedales is compounded by their relative scarcity, there has been a long tradition among Andean communities of enhancing these natural wetlands in order to increase their size and to improve the quality of the pasturage (e.g. Kuznar 1995; Palacios Rios 1996; Carhuallanqui 1998; Quesada & Lema 2011; Verzijl & Guerrero Quispe 2013). As the sixteenth-century Indigenous chronicler Felipe Guaman Poma de Ayala

(1993: 780) notes in his admonition to King Philip III of Spain, indigenous water management systems needed to be maintained—systems that canalised “water for irrigating said fields, even the pasture of the herds they irrigated in the highlands and in the valleys” (authors’ translation).

Archaeologically, these systems have been difficult to detect, mainly due to the often-ephemeral nature of the technologies used to irrigate these pastures artificially, such as the use of unlined canals cut into the fresh sod, and also as a result of continued siltation. Nonetheless, the use of such systems to enhance Andean wetlands has been suggested to have commenced from at least the Middle Horizon (AD 600–1000) onwards and has been linked to the rise of the Tiwanaku state in that period (Vining & Williams 2020: 13). Domic and colleagues (2018: 13) have sought to push back the start of anthropogenic wetland intervention in northern Chile even further, to 2000 years ago.

Among the varied types of infrastructure, Lane (2009: 177–78) identifies silt reservoirs—principally composed of low walls and sluices—as a particular pre-Hispanic form of terrace analogous to check dams or cross-channel terraces documented in the general literature (Denevan 2001: 75) but with one key difference. Check dams and cross-channel terraces are usually found in the middle or lower reaches of side valleys and ravines to counteract the effects of erosion, as well as regulating the flow of water and trapping silt. Check dams of this type were first documented by Regal Matienzo (2005: 72) in the Río Chira Basin in 1970, while other instances were noted by Donkin (1979) in Ecuador, Chile and north-west Argentina. In the latter region, for example, a recent study by Zuccarelli Freire and colleagues (2021) highlights the use of these features across narrow gullies to counteract the effects of land erosion.

Although the silt reservoirs we have previously identified in the Ancash highlands are similar types of structure, they are located at a much higher altitude—generally at the upmost level of a drainage, or along its upper reaches (Lane 2009). The altitude at which these features are found underlines their basic function; aside from controlling water and sediment, they provide the technical means through which to enlarge or create artificial wetlands with enriched pasturage for domesticated animals—in this case, llamas and alpacas.

That these features are also recorded at much lower elevations, within the upper limits of modern-day agriculture, demonstrates that the range of high-altitude Andean pastoralism was far wider in the past than today (Lane & Grant 2016). Documented in the north-central Andean highlands, as well as in the Socaire region of the Chilean Altiplano (Donkin 1979: 126; Lane 2009: 177–78), these features are located in the *sumi* (transitional farming-herding zone: 3500–4100m asl) and puna (alpine-tundra herding grassland: 4100–5000m asl) ecozones of the Andes. Here, we analyse analogous silt reservoirs discovered in the Marccaranca bofedal in the upper Ica drainage of the south-central Andes. As with the north-central Andean examples, the defining feature of these structures is their capacity to store water, manage sediments and generate a micro-environment in which improved pasturage provided the means for successful camelid herding.

SAN 3: the Marccaranca bofedal

The Marccaranca bofedal (13°46′12.9″S, 75°12′36.2″W) is located within the puna ecozone, at approximately 4050m asl. At the top of the eponymous side valley and stream—a

headwater tributary of the Ica River—the bofedal is found within prime herding territory, as attested by the presence of numerous abandoned corrals (most likely of pre-Hispanic and early modern date) (Figures 1 & 2). In its present state, the bofedal covers an area of 8ha. Today there are still two sizable alpaca herds (>250 head) based in the vicinity.

The natural Marccaranca bofedal has been augmented by six silt reservoirs (A–F), covering two spring gullies (A–B & C), a central area (D–E), and, finally, the bofedal stream outlet (F), downslope through the Marccaranca Valley (see Figure 3). The springs feeding into silt reservoirs A–B and C emerge from the Incahuasi granodiorite bedrock uplands that form the prevailing geology at this location. The underlying geology of the bofedal and the area of Incahuasi granodiorite is a phaneritic-textured, intrusive, igneous rock that is similar to granite but containing more plagioclase than orthoclase feldspar (Palacios Moncayo 1994).

Incahuasi granodiorite has a low relative porosity of 0.55–0.59 per cent, with a crack porosity similar to that of Andesite (10–20 per cent) (Schild *et al.* 2001: 118). Thus, in any bofedal basin it underlies, water tends to pool while still seeping through natural cracks, replenishing underground aquifers and springs located downslope. It is therefore likely that this bofedal was altered for the dual purposes of expanding wetland pastures and providing an increase in water for geological storage and subsequent use in terrace agriculture downslope.

In general, while accumulating peat has buried large parts of the silt reservoir structures, in some areas, it is possible to see that the original walls would have been double-faced, infilled with clay and smaller stones, with a maximum thickness of between 1.2 and 1.4m. The reservoir walls, built on the then-existing ground surface, are composed of irregular stone coursing, with larger foundational stones at the base and middling to smaller stones towards the top. The coursing was held together by a compact, silty-clay mortar. In many places, the upper sections of the reservoir have collapsed, leaving only the large base stones visible above ground. The maximum height above the ground surface of these surviving structures is 1.4m, although in most cases, only approximately 0.5–0.7m is currently preserved (Figure 4). A test-pit excavation near to profile 2 (see Figure 5) revealed that silt reservoir E's wall reached a depth of 0.7m, providing an indication of the depth of siltation since construction—at least potentially for those located in the bofedal basin (D–F). The wetness of the reservoirs' exterior walls demonstrates that, while maintaining wet conditions behind them, excess water would have seeped through.

Silt reservoirs A, B and C are located across the two small spring ravines that feed into the main bofedal basin (Figure 6). Silt reservoirs D and E are found below A, B and C to capture water and sediments from the western side of the bofedal. Silt reservoir E abuts D, suggesting that it was built later, although the time elapsed between their construction is unknown. Finally, silt reservoir F is positioned at the lower southern edge of the bofedal, prior to it entering the upper section of the Marccaranca Valley. Silt reservoir F is, therefore, foundational in shoring up the bofedal and facilitating water and soil storage in the larger bofedal basin. All the silt reservoirs utilise existing freestanding Incahuasi granodiorite blocks where present, with the northern end of D, E and F anchored directly onto outcrops of the same rock, lending these structures additional support. The end sections of silt reservoirs A, B and C are freestanding.




Figure 1. Map of Peru, showing the location of the study area (figure by L. Coll).




Figure 2. Map of the study area, showing the Chocorvos-Inca-Spanish Colonial settlement site of Viejo Sangayaico (SAN 1a-e), bofedal of Marccaranca (SAN 3) and pre-Hispanic ritual platform (ushnu), and the present-day calvary of Huinchucruz (SAN 4) (figure by L. Coll).

Discussion: peat, pollen and dates

During fieldwork conducted in 2013, three borehole profiles were taken from the Marccaranca bofedal (Figure 5). Profiles 2 and 3 were located behind silt reservoir E, with profile 2 flush against its inner wall and profile 3 located 5m to the north-west of profile 2. Profile 6 was sited approximately 50m from profile 2, in the middle of the bofedal, and some 75m north of silt reservoir F. Two radiocarbon dates were obtained for each borehole profile (Table 1). In 2018, a small test-pit was dug alongside the profile 2 borehole to ascertain the depth of the wall at this point. A radiocarbon date from the base of this wall was then used to estimate the probable construction date of silt reservoir E.

The data from all three boreholes confirm the anthropogenic manipulation of the Marccaranca bofedal starting in the pre-Hispanic period. Profile 6, moreover, is particularly significant, as it records a sequence of five distinct stages of change in the composition of the Marccaranca bofedal basin over an even longer period of time (Figure 6): 1) an initial humic and peat deposition episode; 2) humic lenses and gritty silting; 3) a black peat episode; 4) a grey silt of a slowly in-filling lake; 5) a final episode of peat formation and gritty lens, creating the bofedal we see today.

The two radiocarbon dates from profile 6, obtained at depths of between 1.57–1.60m and 2.16–2.20m, date the lower section of this profile to 5240 cal BC (5320–5060 BC, 95%




Figure 3. The Marccaranca bofedal (SAN 3), showing silt reservoirs A–E (photograph by K. Lane).



Figure 4. Detail of silt reservoir E, with Sturt Fraser, showing maximum preserved wall height (photograph by K. Lane).




Figure 5. Drone flight panoramic and plan of SAN 3, with profile sediment descriptions (figure by L. Coll).



Figure 6. Detail of silt reservoir C, showing the extent of the bofedal area behind the reservoir wall (photograph by K. Lane).

probability; OxA-32393) and 5680 cal BC (5740–5620 BC, 95% probability; OxA-32394), respectively (Table 1). These dates record, respectively, the episode of black peat (Stage 3) and the basin's initial peat level (Stage 1), demonstrating that bofedal-type conditions have persisted here for the last seven millennia. Later episodes of silting evident in the profile demonstrate the region's dynamic environment. It is possible that the slow infilling lacustrine environment represented by Stage 4 was caused by construction of the silt reservoirs, thus leading to the retention of greater volumes of water in this part of the basin. Subsequent peat formation reveals the stabilisation of the basin's ecological system, leading to its present-day state.

Profiles 2 and 3 (Figure 6), located on the edge of the bofedal, are much more homogeneous than profile 6, showing base levels of fine gravel (like profile 6), followed by long sequences of brown or black peat up to the present day. Figure 7 shows that the deepest parts of profiles 2 and 3 have median radiocarbon dates of 1420 cal BC (1500–1300 BC, 95% probability; OxA-32498) and cal AD 740 (AD 670–880, 95% probability; OxA-32392), respectively (Table 1). These pre-date the creation of silt reservoir E and demonstrate that wetland conditions here persisted through the Late Holocene up to the end of the first millennium AD.

The latter two samples from profiles 2 and 3 date the use of the bofedal after the building of silt reservoir E. The profile 3 date has a median of AD 1850 (AD 1690–1930, 95% probability; OxA-32352), showing continued peat formation from the eighteenth to the twentieth century (Figure 7). The median of the profile 2 date is cal AD 1440 (1410–1460, 95%

Table 1. Radiocarbon dates from borehole samples from the Marccarana bofedal. All samples are peat. $\delta^{13}\text{C}$ estimated by IRMS. Dates calibrated with OxCal v4.4.4 (Bronk Ramsey 2021) using a mixed calibration curve (Marsh *et al.* 2018) of SHCal20 and IntCal20 (Reimer *et al.* 2020). Medians and ranges rounded by 10 years.

Lab code	Profile (sample code)	Depth (m)	$\delta^{13}\text{C}$ ‰	^{14}C age BP	±	Median	95% probability range	Material
OxA-32351	Profile 2 (312)	0.65–0.68	–26.06	492	24	AD 1440	AD 1410–1460	Bulk peat
OxA-32498	Profile 2 (309)	2.04–2.06	–24.49	3168	30	1420 BC	1500–1300 BC	Bulk peat
OxA-32352	Profile 3 (314)	0.60–0.64	–25.00	51	25	AD 1850	AD 1690–1930	Bulk peat
OxA-32392	Profile 3 (315)	1.71–1.73	–26.28	1273	26	AD 740	AD 670–880	Bulk peat
OxA-32393	Profile 6 (299)	1.57–1.60	–27.04	6267	35	5240 BC	5320–5060 BC	Bulk peat
OxA-32394	Profile 6 (300)	2.16–2.20	–26.09	6803	39	5680 BC	5740–5620 BC	Bulk peat




Figure 7. Top) drone flight panoramic and plan of SAN 3, indicating the position of silt reservoirs and profiles; bottom) section of profile locations and radiocarbon samples (figure by L. Coll).

probability; OxA-32351), which comes from the base of the silt reservoir E wall and provides a critical *terminus ante quem* date range for the building of this structure within the first half of the fifteenth century—sometime during the early Inca period.

Recent scholarship has endeavoured to show that the traditional document-based Inca chronology (Rowe 1945) does not reflect the actual time depth of imperial expansion

(e.g. Ogburn 2012; Marsh *et al.* 2017). Indeed, recent dates from the nearby southern coast Acari Valley suggests an Inca presence as early as AD 1430–1460 (Valdez *et al.* 2020: 1569), significantly earlier than Rowe's ethnohistorically informed AD 1476 date (see also Bongers *et al.* 2018: 227 for the mid-Chincha Valley immediately to the north). These studies suggest Inca expansion into the region by the early to mid fifteenth century, which agrees with our profile 2 date that marks the construction of at least some of the bofedal infrastructure at Marccaranca. This is key, given the wider evidence for Inca investment in hydraulic infrastructure across the Andes (e.g. Regal Matienzo 2005). The possibility of imperially driven construction at the Marccaranca bofedal is lent further support by the proximity of the nearby Inca Road and the important Inca occupation at Viejo Sangayaico (SAN 1), located just 3km immediately downslope in an area of extensive bench terrace construction (Lane *et al.* 2016).

Pollen samples were also taken at 40mm intervals from all three profile cores. Selected subsamples were processed using standard techniques for the extraction of sub-fossil pollen and spores (Moore *et al.* 1999). Samples of 1.5ml were analysed and a sum of 300–400 pollen grains plus spores was counted for each sample. Pollen flora were identified using published works on regional pollen and a comparative collection from the Palaeoecology Laboratory at the University of Southampton, UK. The pollen data are plotted using Tilia and Tilia View (TiliaIT 2021), with pollen calculated as a percentage of the total pollen and spores as a percentage of the sum + spores (Figures 8–10). The few tree/shrub taxa are plotted as a group (left of each diagram), while the dominant herb flora are plotted in alphabetic order of family.

All three profiles show clear evidence of bofedal-type plant communities—dominated by plants from the Poaceae, *Plantago*, Cyperaceae and Asteraceae families—persisting from the Middle Holocene (profile 6) through to the present day (profiles 2 and 3). Because of their relative proximity, profiles 2 and 3 show clear correlations; for instance, in their high counts of Poaceae and Asteraceae (the latter is documented as *Tubuliflorae* indet. in the profile pollen diagrams), between their stages 2 and 3, respectively. The upper levels of profile 2, stage 3, however, contain high counts of Cyperaceae and *Plantago*, which are apparently not reflected in the uppermost levels of profile 3, stage 4, probably due to local variations in the bofedal's vegetation.

Strikingly, all the profiles show evidence for maize (*Zea* sp.) pollen in pre-Hispanic contexts. Self-pollinating maize produces a very low pollen rain of heavy grains, which is never dispersed far from the plant. Its presence here therefore suggests its import from lower altitudes for consumption by herders, for foddering animals (as is the case today; McCorkle 1987), or perhaps even high-altitude cultivation. The cultivation of maize pollen could be possible in unused corrals using guano-enriched soils, with the corral walls and the proximity of the bofedal providing a buffer against the wind and cold. Similar shared use of space for herding and farming is recorded throughout the Andes, including cultivation of maize and potato in the seasonal *qochas* or ponds of the circum-Titicaca Basin at nearly 4000m asl (Flores Ochoa *et al.* 1996; Craig *et al.* 2011).

While bofedal-type grasses, sedges and herbs, such as *Plantago* and Asteraceae, dominate these pollen sequences, in profile 2 there is a marked decline in *Plantago* and wetland sedges (Cyperaceae) that is accompanied by a corresponding peak in grasses (Poaceae). This change occurred after the silt reservoir's construction and is dated to cal AD 1440 (AD 1410–1460,




Figure 9. Profile 3 pollen diagram, with calibrated radiocarbon medians. Dates calibrated with OxCal v4.4.4 (Bronk Ramsey 2021) using a mixed calibration curve (Marsh et al. 2018) of SHCal20 and IntCal20 (Reimer et al. 2020) (figure R. Scaife).




Figure 10. Profile 6 pollen diagram, with calibrated radiocarbon medians. Dates calibrated with OxCal v4.4.4 (Bronk Ramsey 2021) using a mixed calibration curve (Marsh et al. 2018) of SHCal20 and IntCal20 (Reimer et al. 2020) (figure by R. Scaife).

95% probability; OxA-32351; see [Figure 8](#)). This sequence is precisely what one would expect, given that increased grass production would have enhanced the herding potential of the bofedal. Such bofedal enhancement under Inca auspices has been documented for the Inca period (Chepstow-Lusty *et al.* 2009). Subsequently, grasses decline, and sedges increase (see Chepstow-Lusty *et al.* 2009: 779–80), presumably as a result of the system falling into disuse following a catastrophic collapse in camelid numbers triggered by the introduction of Old World animals and animal diseases during the early Spanish colony (Flores Ochoa 1980). Profile 3, however, appears to be comparable with a lower-resolution chronology (see [Figure 9](#)).

Conclusions

The Marccaranca bofedal is one component of a landscape transformed through time, first by the native Chocorvos, a Late Intermediate Period (AD 1000–1400) ethnic group (Rowe 1946: 188), and then by the Inca (AD 1400–1532). Our data suggest that at least one, and therefore possibly some of the other silt reservoirs at Marccaranca, date to the early Inca period—tracking other imperial developments in the basin, including the occupation of Sector B at Viejo Sangayaico (SAN 1 in [Figure 2](#); Lane *et al.* 2016), which is located only 3km downslope and which has extensive associated agricultural terracing (Nanavati *et al.* 2016).

At 3650m asl, the multi-period site of Viejo Sangayaico is located precisely at the transition between the agricultural suni and herding puna ecozones, thereby providing the setting for multiple social, economic and religious interactions between their respective herder and farmer communities (*sensu* Parsons *et al.* 1997). Once incorporated into the empire, Viejo Sangayaico was the main Chocorvos and Inca settlement, controlling south–north access along the section of Inca Road that ran adjacent to the Marccaranca bofedal and linked the upper Ica drainage to the northern Pisco drainage. The latter is a particularly important transit route that connected the highlands and the coast with major road and imperial infrastructure, including the administrative centres at Tambo Colorado and Huaytará (Bueno Mendoza 2003; Serrudo Torobeo 2010).

The Marccaranca bofedal thus represents a key element within this wider region of rich ecological resource zones, which included agricultural terraces, along with natural bofedales and pastures—all providing surplus and supplies for the imperial requirements in the Pisco drainage and beyond. This interpretation is supported by ethnohistorical documentation of the Chocorvos shifting their powerbase from the upper Ica to the Pisco drainage, in accordance with the particularly strong Inca state presence there (Espinoza Soriano 2019: 993–1000).

The Marccaranca bofedal's pollen record provides another line of evidence for how imperial investment in this landscape expanded pasture. The presence of maize pollen here at altitudes above 4000m asl hints at attempts to intensify production of this important pre-Hispanic staple crop by pushing its extreme ecological limits (Staller 2016).

A neo-Wittfoglian position, while questioning the primacy of the state in initiating large hydraulic engineering projects, nevertheless advocates the subsequent intervention of the state in community-based systems geared towards increased production (cf. Wittfogel

1957; see Stanish 1994; Lane 2009). Here, we propose just such a model, whereby Chocorvos productive resources, such as terraces, canals and bofedales, were systematised and enlarged via Inca imperial intervention across the entire continuum of hydrological manipulation. In effect, Inca hegemony in the upper Ica drainage seems to have been directed towards a politico-economic objective, attained through engineering and infrastructure investment in settlements, terraces, roads and pasture, and aimed towards the satisfaction of the imperial body-politic, which is especially evident throughout the all-important Pisco drainage that connected the Empire's highland heartland to its coastal hinterlands.

Elsewhere in the Americas, similar state or imperial intervention and enlargement of community-based systems have been suggested for the highland Middle Horizon (AD 600–1000) Tiwanaku polity and the coastal Late Intermediate Period (AD 1000–1400) Chimú state (Graffam 1992; Orloff 1993). Further afield in Mesoamerica, Mayan (400 BC–AD 1000) state action has also been posited as complementing local forms of water management (Wyatt 2014). Indeed, the incidence of top-down intervention over local water-management systems seems to be a constant across the world, especially when the area in question was critical to a state or imperial polity. In this regard, the Roman Empire and its successor state in the east, Byzantium (27 BC–AD 1453), understood the importance of harnessing water as both a resource and a politico-ideological tool to further propagate the state (Wilson 2012).

Likewise, this has been observed in ancient India, China and elsewhere (e.g. Wu *et al.* 2019; Kumar Singh *et al.* 2020), suggesting that while Wittfogel (1957) was incorrect in his contention that water infrastructure investment was the sole prerogative of ancient states and empires (see also Steward 1955), he was correct in gauging the capacity for powerful polities to restructure and increase productivity through recourse to hydraulic engineering, leading to an expansion in state outreach and control. Similarly, in the present day, there seems to be a return to big-state investment in hydraulic engineering, this time more closely tied to local initiatives, sometimes as a means to counter the ever-increasing threat of climate change (Crow-Miller *et al.* 2017). It would therefore seem useful for archaeologists to revisit the wider implications of Wittfogelian theory concerning the primacy of water management for ancient (and not so ancient) polities, endeavouring, in each case, to disentangle the political ecology underlying state intervention, regardless of whether such intervention was ultimately successful or not.

Acknowledgements

We thank all the members of the *Proyecto de Investigación Arqueológica Cuenca de Ica* (PIACI) and *One River Project*, as well as the local community of Sangayaico for all their help in making this project possible. Permission for the project was granted by the Ministerio de Cultura del Perú under RD nos 061-2013-DGPA-VMPCIC/MC and 430-2018-DGPA-VMPCIC/MC.

Funding statement

Funding for the project was obtained from the Leverhulme Foundation (UK [Project Code: RPG-17]: 2012–2016) and the Gerda Henkel Foundation (Germany [Project Code: AZ-30-F-17]: 2018–2021).

References

- BONGERS, J.L., C. O'SHEA & A. FARAHANI. 2018. Communities of weavers: a methodology for analyzing textile and cloth production. *Journal of Archaeological Science Reports* 22: 223–36. <https://doi.org/10.1016/j.jasrep.2018.09.025>
- BRONK RAMSEY, C. 2021. OxCal v4.4 manual. Available at: http://intchron.org/tools/oxcalhelp/hlp_contents.html#Introduction (accessed 5 December 2021).
- BROWMAN, D.L. 1990. High altitude camelid pastoralism of the Andes, in J.G. Galaty & D.L. Johnson (ed.) *The world of pastoralism: herding systems in comparative perspective*: 323–52. New York: Guilford Press.
- BUENO MENDOZA, A. 2003. El Tiwanintinsuyu en Huaytará. *Investigaciones Sociales* 7: 41–56. <https://doi.org/10.15381/is.v7i11.8103>
- CARHUALLANQUI, R.M. 1998. *Pastores de Altura: Magia, Ritos y Danzas*. Lima: REDES: Red de Solidaridad.
- CHEPSTOW-LUSTY, A.J. *et al.* 2009. Putting the rise of the Inca Empire within a climatic and land management context. *Climate of the Past* 5: 771–96. <https://doi.org/10.5194/cpd-5-771-2009>
- CRAIG, N. *et al.* 2011. Geologic constraints on rain-fed Qocha reservoir agricultural infrastructure, northern Lake Titicaca basin, Peru. *Journal of Archaeological Science* 38: 2897–907. <https://doi.org/10.1016/j.jas.2011.05.005>
- CROW-MILLER, B., M. WEBBER & F. MOLLE. 2017. The (re)turn to infrastructure for water management? *Water Alternatives* 10: 195–207.
- DENEVAN, W.M. 1992. The pristine myth: the landscape of the Americas in 1492. *Annals of the Association of American Geographers* 82: 369–85. <https://doi.org/10.1111/j.1467-8306.1992.tb01965.x>
- 2001. *Cultivated landscapes of native Amazonia and the Andes*. Oxford: Oxford University Press.
- DOMIC, A.I. *et al.* 2018. Two thousand years of land-use and vegetation evolution in the Andean highlands of northern Chile inferred from pollen and charcoal analyses. *Quaternary* 1: 32. <https://doi.org/10.3390/quat1030032>
- DONKIN, R.A. 1979. *Agricultural terracing in the aboriginal New World* (Viking Fund Publications in Anthropology 56). Tucson: University of Arizona Press.
- ELLIS, E.C. *et al.* 2021. People have shaped most of terrestrial nature for at least 12 000 years. *Proceedings of the National Academy of Sciences of the USA* 118: e2023483118. <https://doi.org/10.1073/pnas.2023483118>
- ESPINOZA SORIANO, W. 2019. *Etnias del Imperio de los Incas: Reinos, señoríos, Curacazgos y Cacicatos (Volumen II)*. Lima: Universidad Ricardo Palma Editorial Universitaria.
- FLORES OCHOA, J.A. 1980. Causas que originaron la actual distribución de las alpacas y llamas, in L. Millones & H. Tomeada (ed.) *El Hombre y su Ambiente en los Andes Centrales* (Senri Ethnological Studies 10): 63–92. Osaka: National Museum of Ethnology.
- FLORES OCHOA, J.A., M.P. PAZ FLORES & W. ROZAS. 1996. Un (re)-descubrimiento reciente: la agricultura en lagunas temporales (qocha) en el Altiplano, in P. Morlon (ed.) *Comprender la agricultura campesina en los Andes Centrales: Perú-Bolivia*: 247–55. Lima: IFEA/CBC.
- GRAFFAM, G. 1992. Beyond state collapse: rural history, raised fields, and pastoralism in the south Andes. *American Anthropologist* 94: 882–904. <https://doi.org/10.1525/aa.1992.94.4.02a00060>
- KUMAR SINGH, P., P. DEY, S.K. JAINA & P.P. MUJUMDAR. 2020. Hydrology and water resources management in ancient India. *Hydrology and Earth System Sciences* 24: 4691–707. <https://doi.org/10.5194/hess-24-4691-2020>
- KUZNAR, L.A. 1995. *Awatimarka: the ethnology of an Andean herding community*. Fort Worth (TX): Harcourt Brace College.
- LANE, K. 2006. Through the looking glass: re-assessing the role of agro-pastoralism in the north-central Andean highlands. *World Archaeology* 38: 493–510. <https://doi.org/10.1080/00438240600813806>
- 2009. Engineered highlands: the social organisation of water in the ancient north-central Andes (AD 1000–1480). *World Archaeology* 41: 169–90. <https://doi.org/10.1080/00438240802655245>
- LANE, K. & J.L. GRANT. 2016. A question of altitude: exploring the limits of highland pastoralism in the Prehispanic Andes, in J.M. Capriles & N. Tripcevich (ed.) *The archaeology of Andean pastoralism*: 139–57. Albuquerque: University of New Mexico Press.

- LANE, K. *et al.* 2016. Anatomía de un Abandono: El caso de Viejo Sangayaico [SAN 1]. *Inka Llaqta* 4: 85–98.
- MALDONADO FONKÉN, M.S. 2014. An introduction to the *bofedales* of the Peruvian high Andes. *Mires and Peat* 15: 1–13.
- MARSH, E.J., R. KIDD, D. OGBURN & V. DURÁN. 2017. Dating the expansion of the Inca Empire: Bayesian models from Ecuador and Argentina. *Radiocarbon* 59: 117–40. <https://doi.org/10.1017/RDC.2016.118>
- MARSH, E.J. *et al.* 2018. IntCal, SHCal, or a mixed curve? Choosing a ¹⁴C calibration curve for archaeological and paleoenvironmental records from tropical South America. *Radiocarbon* 60: 925–40. <https://doi.org/10.1017/RDC.2018.16>
- MCCORKLE, C.M. 1987. Punas, pastures and fields: grazing strategies and the agropastoral dialectic in an Indigenous Andean community, in D.L. Browman (ed.) *Arid land use strategies and risk management in the Andes: a regional anthropological perspective*: 57–80. Boulder (CO): Westview.
- MOORE, P.D., J.A. WEBB & M.E. COLLINSON. 1999. *Pollen analysis*. Oxford: Blackwell Science.
- NANAVATI, W.P. *et al.* 2016. Testing soil fertility of Prehispanic terraces at Viejo Sangayaico in the upper Ica catchment of south-central highland Peru. *Catena* 142: 139–52. <https://doi.org/10.1016/j.catena.2016.03.007>
- OGBURN, D.E. 2012. Reconceiving the chronology of the Inca imperial expansion. *Radiocarbon* 54: 219–37. https://doi.org/10.2458/azu_js_rc.v54i2.16014
- ORTLOFF, C.R. 1993. Chimu hydraulic technology and statecraft on the north coast of Peru, AD 1000–1470, in V. Scarborough & B. Isaacs (ed.) *Economic aspects of water management in the Prehispanic New World* (Research in Economic Anthropology, Supplement 7): 327–67. Greenwich (CT): JAI Press.
- PALACIOS MONCAYO, O. 1994. *Geología de los Cuadrángulos de Santiago de Chocorvos y Paras, Hojas: 28-m y 28-n* (Serie A: Carta Geológica Nacional Boletín 49). Lima: Instituto Geológico Minero y Metalúrgico, Sector Energía y Minas.
- PALACIOS RIOS, F. 1996. Pastizales de regadío para alpacas en la puna alta (El ejemplo de Chichillapi), in P. Morlon (ed.) *Comprender la agricultura campesina en los Andes Centrales: Perú-Bolivia*: 207–12. Lima: IFEA/CBC.
- PARSONS, J.R., C.M. HASTINGS & R. MATOS MENDIETA. 1997. Rebuilding the state in highland Peru: herder-cultivator interaction during the Late Intermediate Period in the Tarama-Chinchaycocha region. *Latin American Antiquity* 8: 317–41. <https://doi.org/10.2307/1972106>
- POMA DE AYALA, F.G. 1993. *Nueva Crónica y Buen Gobierno* (Volumes 1–3). Translated by F.G.Y. Pease. Lima: Fondo de Cultura Económica.
- QUESADA, M.N. & C. LEMA. 2011. Los Potreritos de Antofagasta: trabajo Indígena y Propiedad (finales del Siglo XVIII y comienzos del XIX). *Andes* 22: 1–21.
- RAVINES, R. 1978. *Tecnología Andina* (Fuentes e Investigaciones para la Historia del Perú 4). Lima: IEP/ITINTEC.
- REGAL MATIENZO, A. 2005. *Los trabajos hidráulicos del inca en el antiguo Perú*. Lima: Instituto Nacional de Cultura.
- REIMER, P.J. *et al.* 2020. The IntCal20 Northern Hemisphere radiocarbon age calibration curve (0–55 cal kBP). *Radiocarbon* 62: 725–57. <https://doi.org/10.1017/RDC.2020.41>
- ROWE, J.H. 1945. Absolute chronology in the Andean area. *American Antiquity* 10: 265–84. <https://doi.org/10.2307/275130>
- 1946. Inca culture at the time of the Spanish conquest, in J.H. Steward (ed.) *Handbook of South American Indians* (Bureau of American Ethnography, Bulletin No. 143): 183–330. Washington, D.C.: Smithsonian Institution.
- SCHILD, M., S. SIEGESMUND, A. VOLLBRECHT & M. MAZUREK. 2001. Characterization of granite matrix porosity and pore-space geometry by *in-situ* and laboratory methods. *Geophysical Journal International* 146: 111–25. <https://doi.org/10.1046/j.0956-540x.2001.01427.x>
- SERRUDO TOROBEJO, E. 2010. El Tampu Real de Inkahuasi y la ocupación Inka en Huaytara. *Inka Llaqta: Revista de Investigaciones Arqueológicas y Etnohistóricas Inka* 1: 173–93.
- SQUEO, F.A., B.G. WARNER, R. ARAVENA & D. ESPINOSA. 2006. Bofedales: high altitude peatlands of the central Andes. *Revista Chilena de Historia Natural* 79: 245–55. <https://doi.org/10.4067/S0716-078X2006000200010>
- STALLER, J.E. 2016. High altitude maize (*Zea mays* L.) cultivation and endemism in the Lake Titicaca basin. *Journal of Botany Research* 1: 8–21. <https://doi.org/10.36959/771/556>

- STANISH, C. 1994. The hydraulic hypothesis revisited: Lake Titicaca basin raised fields in theoretical perspective. *Latin American Antiquity* 5: 312–32.
<https://doi.org/10.2307/971820>
- STEWART, J.H. 1955. *Irrigation civilizations: a comparative study*. Washington, D.C.: Pan American Union.
- TiliaIT. 2021. *Tilia*. Available at: <https://www.tiliait.com> (accessed 5 December 2021).
- VALDEZ, L.M., K.J. BETTCHER & M.N. HUAMANÍ. 2020. Inka llama offerings from Tambo Viejo, Acari Valley, Peru. *Antiquity* 94: 1557–74.
<https://doi.org/10.15184/aqy.2020.183>
- VERZIJL, A. & S. GUERRERO QUISPE. 2013. The system nobody sees: irrigated wetland management and alpaca herding in the Peruvian Andes. *Mountain Research and Development* 33: 280–93.
<https://doi.org/10.1659/MRD-JOURNAL-D-12-00123.1>
- VINING, B. & P.R. WILLIAMS. 2020. Crossing the western Altiplano: the ecological context of Tiwanaku migrations. *Journal of Archaeological Science* 113: 105046.
<https://doi.org/10.1016/j.jas.2019.105046>
- WILSON, A. 2012. Water, power and culture in the Roman and Byzantine worlds: an introduction. *Water History* 4: 1–9.
<https://doi.org/10.1007/s12685-012-0050-2>
- WITTFOGEL, K.A. 1957. *Oriental despotism: a comparative study of total power*. New Haven (CT): Yale University Press.
- WU, S. *et al.* 2019. The development of ancient Chinese agricultural and water technology from 8000 BC to 1911 AD. *Palgrave Communications* 5: 77. <https://doi.org/10.1057/s41599-019-0282-1>
- WYATT, A.R. 2014. The scale and organization of ancient Maya water management. *WIREs Water* 1: 449–67. <https://doi.org/10.1002/wat2.1042>
- YOUNG, K.R., B. LEÓN & O. HERRERA-MACBRYDE. 1997. Peruvian puna, in S.D. Davis *et al.* (ed.) *Centres of plant diversity: a guide and strategy for their conservation. Volume 3: the Americas*: 470–76. Cambridge: WWF/IUCN.
- ZUCCARELLI FREIRE, V., A.S. MELÉNDEZ, M. RODRIGUEZ OVIEDO & M.N. QUESADA. 2021. Erosion control in Prehispanic agrarian landscapes from Northwestern Argentina: El Alto-Ancasti Highlands case study (Catamarca, Argentina). *Geoarchaeology* 37: 74–95.
<https://doi.org/10.1002/gea.21862>